

Mott Insulators Without Symmetry Breaking

Dung-Hai Lee^{a,b}, and Jon Magne Leinaas^{a,c}

(a) *Department of Physics, University of California at Berkeley, Berkeley, CA 94720, USA*

(b) *Material Science Division, Lawrence Berkeley National Laboratory and*

(c) *Department of Physics, University of Oslo, P.O. Box 1048 Blindern, 0316 Oslo, Norway*

(Dated: February 2, 2008)

We present theoretical models, in one and two space dimensions, that exhibit Mott insulating ground states at fractional occupations without any symmetry breaking. The Hamiltonians of these models are non-local in configuration space, but local in phase space.

In recent years the study of doped Mott insulators [1] has become one of the main themes of condensed matter physics [2]. In a Mott insulator the motion of each particle is hindered by the presence of other particles due to their strong repulsive interaction, and in this respect they are different from band insulators where the Pauli exclusion principle forbids the motion of electrons. An interesting feature of Mott insulators is that doping, *i.e.*, changing of particle density, can dramatically affect their properties. For example, doping of “cuprates” and “manginites” has lead to the high transition temperature superconductors and the colossal magneto-resistive materials.

If a crystalline solid has a fractional occupation number [3] and is Mott insulating, there seems always to be a spontaneous symmetry breaking of the translation symmetry [4]. After the symmetry breaking the unit cell is effectively enlarged so that the occupation number becomes an integer. This kind of symmetry breaking blurs the distinction between Mott insulators and band insulators[5].

Nearly thirty years ago, Anderson and Fazekas [6] postulated that due to the frustrated antiferromagnetic interaction, the Mott state on triangular lattice does not break the spin rotation symmetry (hence is “featureless”). This novel Mott state has since been dubbed the name “spin liquid”. Soon after the discovery of high temperature superconductivity, Anderson proposed that the parent (Mott insulating) compounds of these superconductors are spin liquids [7]. Although subsequent developments have shown this is not true, nonetheless this proposal has stimulated wide interest in spin liquids[8].

One of the reasons that we are interested in “featureless” Mott insulators is because it is widely believed that the elementary excitations in such systems may carry fractional quantum numbers (e.g., fractional charge, fractional spin, etc). As a result, the conducting state(s) produced by doping them can be very different from the those produced by doping ordinary symmetry-breaking Mott insulators.

In this paper we present two examples that exhibit featureless Mott insulating ground states. In both cases the elementary excitations carry fractional quantum numbers. From these examples we learn a valuable



FIG. 1: The effect of H_{1D} is to hop a pair of particles while conserving their center of mass position. The hopping range is set by κ^{-1} . In this figure the nearest neighbor pair hopping is illustrated.

lesson in constructing models of this kind.

Without further delay let us define these models. First we focus on a one-dimensional (1D) model. Consider a finite lattice with $M = 3N - 2$ sites where N is an odd integer. We label each site by an integer n with the central site $n = 0$. On this lattice we put N spinless fermions. The occupation number is given by $\nu = N/(3N - 2)$ which approaches $1/3$ in the $N \rightarrow \infty$ limit. It turns out that this choice of the number of sites ensures that the ground state is non-degenerate for all N . The Hamiltonian that describes the interaction between these N fermions is given by

$$H_{1D} = g\kappa^3 \sum_p b_p^\dagger b_p. \quad (1)$$

In the above g is an energy parameter, and is positive. The parameter κ , whose meaning shall be discussed later, is dimensionless ($0 \leq \kappa < 1$). Since H_{1D} is a sum of positive-definite operators, its eigenvalues are clearly non-negative.

In H_{1D} the index p runs through both integers and half integers, and

$$b_p = \sum_q \left(q e^{-\kappa^2 q^2} \right) c_{p-q} c_{p+q}, \quad (2)$$

where q takes integer/half-integer value depending on whether p is integer/half-integer. In Eq. (2) $c_{p \pm q}$ annihilates a fermion at lattice site $p \pm q$. H_{1D} describes the hopping of pairs of fermions from sites $p - q$ and $p + q$ to $p - k$ and $p + k$ (Fig.1). In such a hopping process the center-of-mass position (p) of the fermion pair remains unchanged. From Eq. (2) it is apparent that κ^{-1} controls the hopping range.

For occupation $\nu = N/(3N - 2)$ and *any* κ , the model

defined by Eq. (1) possesses a ground state given by

$$|\Psi_{1D}\rangle = \sum_{\{n_k\}} \chi(n_1, \dots, n_N) c_{n_1}^+ \dots c_{n_N}^+ |0\rangle, \quad (3)$$

where

$$\begin{aligned} \chi(\{n_j\}) &\propto \prod_j e^{-\kappa^2 n_j^2 / 2} \int_{-\infty}^{\infty} dx_j \int_0^{2\pi} dy_j e^{-x_j^2 / \kappa^2} \\ &\times e^{n_j(x_j - iy_j)} \prod_{j>k} \sinh^3 \left[\frac{(x_j - x_k) + i(y_j - y_k)}{2} \right]. \end{aligned} \quad (4)$$

This state is annihilated by all the b_p 's, *i.e.*,

$$b_p |\Psi_{1D}\rangle = 0. \quad (5)$$

As a the result $H_{1D}|\Psi_{1D}\rangle = 0$ implying $|\Psi_{1D}\rangle$ is a ground state of H_{1D} . Eq. (5) can be shown to be true by rewriting it in terms of the wavefunction $\chi(\{n_j\})$ and using the following identity[9]

$$\begin{aligned} \int_{-\infty}^{\infty} dx_1 dx_2 \int_0^{2\pi} dy_1 dy_2 \Phi_p^*(z_1, z_2) \prod_{j>k} \sinh^3 \left(\frac{z_j - z_k}{2} \right) \\ \times e^{-\frac{1}{2\kappa^2} \sum_k x_k^2} = 0, \quad \text{for all } p. \end{aligned} \quad (6)$$

The Φ_p in Eq. (6) is given by

$$\begin{aligned} \Phi_p(z_1, z_2) &= \left[e^{-\kappa^2 p^2} e^{p(z_1 + z_2) - \frac{1}{4\kappa^2} (x_1 + x_2)^2} \right] \\ &\times \left[\sum_q q e^{-2\kappa^2 q^2} e^{q(z_1 - z_2) - \frac{1}{4\kappa^2} (x_1 - x_2)^2} \right]. \end{aligned} \quad (7)$$

In latter part of the paper we shall show that for sufficiently small positive values of κ , $|\Psi_{1D}\rangle$ is the unique ground state of H_{1D} , it separated by an energy gap from the lowest excited states, and it does not break translational symmetry. Meanwhile, to get a feeling for $|\Psi_{1D}\rangle$, let us look at a representation of it for 3 particles, as shown in Fig.2. The result is a coherent superposition of many different configurations. The relative weight and phase of these configurations ensures that they are annihilated by b_p . Any breaking of the weight-phase relation causes excited states.

Next we present a two dimensional (2D) model with similar properties. The lattice consists of M' chains, with each chain made up of $M = 3N - 2$ sites. In each chain we place N spinless fermions. As the result, the 2D occupation number is also given by $\nu = N/(3N - 2)$. In the limit $N, M' \rightarrow \infty$ the occupation number is also $1/3$. The Hamiltonian of the 2D model is given by

$$\begin{aligned} H_{2D} &= \sum_{n,p} \left[g_1 \kappa^3 b_{(n,p)}^+ b_{(n,p)} + g_2 \kappa^3 (b_{(n,p)}^+ b_{(n+1,p)} \right. \\ &\quad \left. + b_{(n+1,p)}^+ b_{(n,p)}) \right]. \end{aligned} \quad (8)$$




FIG. 2: $|\psi_{1D}\rangle$ for three particles. It is represented by components with different occupation of sites. The relative weight and phase between these configuration are $e^{9\kappa^2}$ (top), $3e^{7\kappa^2}, 3e^{7\kappa^2}, 6e^{4\kappa^2}, -15e^{\kappa^2}$ (bottom) respectively.

In the above the chain index n runs through integers and p can be both integer and half-integers. In H_{2D}

$$b_{n,p} = \sum_q \left(q e^{-\kappa^2 q^2} \right) c_{(n,p-q)} c_{(n,p+q)}, \quad (9)$$

where q runs through integer/half-integer depending on whether p is integer/half-integer. The energy parameters g_1 and g_2 are both positive. The first term of H_{2D} hops a pair of particles within each chain n , and the second term hops a pair of particles from chain n to chain $n+1$ and vice versa. In both hopping processes the center of mass coordinate along the chain direction remains unchanged.

For any values of g_1 and g_2 ,

$$|\Psi_{2D}\rangle = \prod_n |\Psi_{1D,n}\rangle, \quad (10)$$

where $|\Psi_{1D,n}\rangle$ is the previous $|\Psi_{1D}\rangle$ defined for the n th chain, is a zero-energy *eigen state* of H_{2D} . This is because b_p annihilates $|\Psi_{1D}\rangle$. Let us imagine obtaining H_{2D} by turning on g_2 from zero while holding g_1 fixed. At non-zero g_2 the excited states of adjacent chains are mixed. However, so long as g_2 is small compared to the excitation gap at $g_2 = 0$, this mixing will not produce negative energy eigen states.[10] Under that condition $|\psi_{2D}\rangle$ continues to be the exact ground state of H_{2D} , and it is separated by an energy gap from the excited states. Since $|\psi_{1D}\rangle$ does not break translation symmetry, so does't $|\Psi_{2D}\rangle$.

The remaining task is to collaborate the statements we have made about $|\psi_{1D}\rangle$. To accomplish this we will make use of a mapping from the 1D Hamiltonian to a well-studied problem in two dimensions, namely, the spinless fermions in a strong magnetic field.

It is well known that when the lowest Landau level (LLL) is $1/3$ filled, and if the interaction between fermions falls off sufficiently fast as a function of the interparticle distance, the ground state is an incompressible quantum fluid. It is also well-known that when a particle

is confined to the lowest Landau level the two components of its coordinate no longer commute. This implies that the dynamics of the fermions is effectively that of a 1D system. This intrinsic one-dimensionality can be made explicit through a mapping of the LLL problem to one dimension. What has not been appreciated is that *although the time reversal symmetry is explicitly broken in the 2D problem, the corresponding 1D Hamiltonian is time-reversal invariant*. Moreover, after the mapping to 1D, the ground state is an example of a featureless Mott insulator at $1/3$ occupation number.

To start, let us consider a spin-polarized two dimensional electron gas where the electrons interact through a two-body potential

$$H_{int} = V_0 \int d^2r d^2r' V(\mathbf{r} - \mathbf{r}') \psi^\dagger(\mathbf{r}) \psi^\dagger(\mathbf{r}') \psi(\mathbf{r}') \psi(\mathbf{r})$$

$$V(\mathbf{r}) = \nabla^2 \delta(\mathbf{r}). \quad (11)$$

Although this interaction potential is a singular function of the coordinates, it has non-singular matrix element between states in the lowest Landau level. Laughlin's $\nu = 1/3$ wavefunction [11] is known to be the exact ground state of this potential [12]. The gap to the lowest excited state is proportional to the parameter V_0/l_B^4 , where l_B is the magnetic length.

Next we place the above two dimensional electron gas on a cylinder with circumference L with the magnetic field perpendicular to the surface. In the Landau gauge the LLL basis orbitals are of the following form

$$\phi_n(\mathbf{r}) = \frac{1}{\sqrt{\pi^{1/2} L l_B}} e^{i2n\pi y/L} e^{-(x-2\pi n l_B^2/L)^2/2l_B^2}. \quad (12)$$

These orbitals are delocalized around the cylinder (y) while localized along its axis (x). The quantization of the y -momentum in units of $2\pi/L$ explains why we have a lattice. If we expand the field operator $\psi(\mathbf{r})$, projected to the lowest Landau level, as $\psi(\mathbf{r}) = \sum_n \phi_n(\mathbf{r}) c_n$, substitute the result into Eq. (11), and rescale the coordinates by $L/2\pi$, we obtain the Hamiltonian (1) with $\kappa \equiv 2\pi l_B/L$, and $g = \frac{4V_0}{(2\pi)^{3/2} l_B^4}$. Thus κ measures the ratio between the magnetic length and the circumference of the cylinder. Since κ^{-1} controls the hopping lengths of the 1D and 2D Hamiltonians, it is clearly desirable to restrict $\kappa > 0$. On the other hand, to preserve the the liquid property of the Laughlin state we need to keep $L > l_B$ (or $\kappa < 2\pi$). To satisfy both requirements we assume $0 < \kappa < 1$.

In Landau gauge, the $\nu = 1/3$ Laughlin wavefunction on a cylinder reads [13, 14]

$$\psi_{1/3}(\{x_j, y_j\}) \propto \prod_{j>k} \sinh^3\left(\frac{z_j - z_k}{L/\pi}\right) e^{-\sum_j x_j^2/2l_B^2}, \quad (13)$$

where $z = x + iy$. The $\chi(\{n_k\})$ given in Eq. (4) are the coefficients of $\psi_{1/3}$ when expanded in terms of products of the lowest Landau level orbitals (Eq. (12)).

At this point we have mapped H_{int} to H_{1D} , and the Laughlin state $|\Psi_{1/3}\rangle$ to $|\Psi_{1D}\rangle$. Since the above mapping is a unitary transformation, all known properties of $|\Psi_{1/3}\rangle$ and H_{int} are preserved. For example, the fact that $|\Psi_{1/3}\rangle$ describes a quantum liquid at magnetic filling factor $1/3$ translates into the statement that $|\Psi_{1D}\rangle$ is a quantum liquid at occupation number $1/3$. The fact that $|\Psi_{1/3}\rangle$ is the non-degenerate ground state of H_{int} implies that $|\Psi_{1D}\rangle$ is the non-degenerate ground state of H_{1D} . Finally, the fact that H_{int} possesses an energy gap and fractional charge quasiparticles at filling factor $1/3$ implies the same for H_{1D} at lattice occupation $1/3$. In this way we have established that the model presented by H_{1D} possesses a featureless Mott insulating ground state at fractional occupation number!

By using the 1D to 2D mapping discussed previously it is simple to map H_{2D} to a 3D Hamiltonian H'_{int} describing coupled layers:

$$H'_{int} = \sum_n H_n$$

$$H_n = \int d^2r d^2r' V(\mathbf{r} - \mathbf{r}') \left[V_1 \psi_n^\dagger(\mathbf{r}) \psi_n^\dagger(\mathbf{r}') \psi_n(\mathbf{r}') \psi_n(\mathbf{r}) \right. \\ \left. + V_2 (\psi_{n+1}^\dagger(\mathbf{r}) \psi_{n+1}^\dagger(\mathbf{r}') \psi_n(\mathbf{r}') \psi_n(\mathbf{r}) + h.c.) \right]. \quad (14)$$

In the above equation $V(\mathbf{r})$ is the same as that given in Eq. (11), and $V_{1,2} \propto g_{1,2}$ respectively. With $V_2 = 0$ the problem becomes that of many independent layers. At $\nu = 1/3$ each layer is in the liquid ground state described by $|\Psi_{1/3}\rangle$. The term proportional to V_2 hops a pair of electrons between adjacent layers. As discussed below Eq. (10), so long as V_2 is sufficiently smaller than V_1 , the ground state is the direct product of the $1/3$ Laughlin liquid in each layer.

It is interesting to note that unlike usual Hamiltonians, H_{1D} and H_{2D} do not have the single-fermion hopping term. One can in fact add a single particle hopping term

$$H'_{1D} = -t e^{-\kappa^2/4} \sum_n [c_{n+1}^\dagger c_n + h.c.] \quad (15)$$

to H_{1D} without changing qualitatively any of the results. To understand this we note that after mapping to 2D Eq. (15) becomes

$$-t \int d^2r \cos(2\pi y/L) \psi^\dagger(\mathbf{r}) \psi(\mathbf{r}), \quad (16)$$

i.e., a periodic potential. It is clear that so long as $|t|$ is much smaller than the energy gap of H_{int} , it will not change the quantum liquid nature of the ground state and will not collapse the excitation gap. Similar single-particle terms can be added to H_{2D} . Again, so long as the strength of the single particle hopping terms is sufficiently small, the qualitative results we discussed above will be preserved.

Models that gives featureless Mott insulating ground state at fractional occupation number have been difficult to find. What is special about the models presented in this paper? To answer this question let us focus on H_{1D} . A standard (1D) interaction potential is local in configuration space (q-space), i.e.,

$$\hat{V} = \prod_j \int dq_j \sum_{i < j} U(q_i - q_j) |q_1, \dots, q_N\rangle \langle q_1, \dots, q_N| \quad (17)$$

When this type of potential dominates the Hamiltonian, it favors particles to assume a fixed q-space configuration. For a repulsive potential such a configuration often take the form of a regular lattice. H_{1D} , on the other hand, is non-local in q-space and the particles do not necessarily have to pay a high price in energy to get close. However, there is a hidden locality in H_{1D} which is revealed not in configuration space, but in *phase space*.

The quantum description of phase space dynamics is most conveniently done in terms of coherent states. For a single particle with coordinate q and momentum p , the coherent state $|z\rangle$ satisfies

$$\left[\hat{q} + i \frac{\lambda^2}{\hbar} \hat{p} \right] |z\rangle = z |z\rangle, \quad (18)$$

where $z = x + iy$ and λ is *any* pre-chosen length scale serving to make $\hat{q}$ and $(\lambda^2/\hbar)\hat{p}$ the same dimension. If $|q\rangle$ denotes the position eigenstate, it is simple to show that

$$\langle q | z \rangle \propto e^{iqy/\lambda^2} e^{-(x-q)^2/2\lambda^2}. \quad (19)$$

and this function defines the transformation between the phase space and configuration space descriptions. It is interesting to note if we identify λ with the magnetic length and q/λ^2 with the y-momentum, the functions given in Eq. (19) become precisely the LLL basis orbitals in Eq. (12). Thus a 2D quantum mechanical problem defined in the LLL is completely equivalent to the coherent state representation of a 1D quantum problem.

Let us go back to the non-local Hamiltonian H_{1D} . When expressed in terms of the coherent state basis, it gets a local form

$$H_{1D} = \sum_{i < j} \int \prod_k \frac{d^2 z_k}{2\pi\lambda^2} V(z_i - z_j) |z_1, \dots, z_N\rangle \langle z_1, \dots, z_N| \quad (20)$$

The $V(z_i - z_j)$ in the above equation is the potential given in H_{int} . This potential prevents particles from getting close together in the phase space. However due to the non-commutativity of the phase space coordinates, the particles are not frozen in any particular configuration. We believe the fact that H_{1D} and H_{2D} are linked

to phase-space local potential is the main reason that they have featureless Mott insulating ground states at fractional occupation numbers.

In conclusion, we have constructed 1D and 2D lattice models that exhibit featureless Mott insulating ground state at fractional occupation number. The Hamiltonians of these models have finite range interactions, and respect lattice translation and time reversal symmetries. These Hamiltonians are local in phase space instead of configuration space. We believe phase space local interaction could be a missing key for constructing models of this type of novel insulators.

DHL is supported by DOE grant DE-AC03-76SF00098. JML thanks the Miller Institute for Basic Research in Science for financial support and hospitality during his recent visit. Supports from the Research Council of Norway and the Fulbright foundation are also acknowledged.

-
- [1] N.F. Mott, *Metal-insulator transitions*, Taylor & Francis, London (1974).
 - [2] M. Imada, A. Fujimori, and Y. Tokura, Rev. Mod. Phys. **70**, 1039 (1998).
 - [3] The occupation number is the average particle number *per spin* in each crystalline unit cell.
 - [4] N. Read and S. Sachdev, Phys. Rev. B **42**, 4568 (1990); D-H Lee and R. Shankar, Phys. Rev. Lett., **65**, 1490 (1990); M. Oshikawa, Phys. Rev. Lett. **84**, 1535 (2000).
 - [5] In a band insulator all low-energy one-particle quantum states are filled. Such a system is insulating because of the Pauli exclusion principle. Usually a band insulator has even number of electrons per unit cell.
 - [6] P. Fazekas and P.W. Anderson, Phil. Mag. **30**, 423 (1974).
 - [7] P.W. Anderson, Science, **235**, 1196 (1987).
 - [8] For representative works on this subject see, e.g., G. Baskaran, Z. Zou and P.W. Anderson, Solid State Commun. **63**, 973 (1987); V. Kalmeyer and R.B. Laughlin, Phys. Rev. Lett. **59**, 2095 (1987); D.S. Rokhsar, and S.A. Kivelson, Phys. Rev. Lett. **61**, 2376 (1988); N. Read and S. Sachdev, Phys. Rev. Lett. **62**, 1694 (1989); R. Moessner and S.L. Sondhi, Phys. Rev. Lett. **86**, 1881 (2001); For a recent review and more references see X.-G. Wen, Phys. Rev. B **65**, 165113 (2002).
 - [9] This requires a straightforward but somewhat lengthy proof.
 - [10] We have performed explicit analysis to second order in g_2 to confirm this statement.
 - [11] R.B. Laughlin, Phys. Rev. Lett. **60**, 2677 (1988).
 - [12] S.A. Trugman and S.A. Kivelson, Phys. Rev. B **31** 5280, (1985).
 - [13] E. Westerberg and T.H. Hansson, Phys. Rev. B **47**, 16 554 (1993).
 - [14] H. Azuma and S. Iso, Phys. Lett. B, **331**, 107 (1994).